

# Intermittence and nonlinear parabolic stochastic partial differential equations\*

Mohammud Foondun      Davar Khoshnevisan  
University of Utah

April 16, 2008

## Abstract

We consider nonlinear parabolic SPDEs of the form  $\partial_t u = \mathcal{L}u + \sigma(u)\dot{w}$ , where  $\dot{w}$  denotes space-time white noise,  $\sigma : \mathbf{R} \rightarrow \mathbf{R}$  is [globally] Lipschitz continuous, and  $\mathcal{L}$  is the  $L^2$ -generator of a Lévy process. We present precise criteria for existence as well as uniqueness of solutions. More significantly, we prove that these solutions grow in time with at most a precise exponential rate. We establish also that when  $\sigma$  is globally Lipschitz and asymptotically sublinear, the solution to the nonlinear heat equation is “weakly intermittent,” provided that the symmetrization of  $\mathcal{L}$  is recurrent and the initial data is sufficiently large.

Among other things, our results lead to general formulas for the upper second-moment Liapounov exponent of the parabolic Anderson model for  $\mathcal{L}$  in dimension  $(1 + 1)$ . When  $\mathcal{L} = \kappa \partial_{xx}$  for  $\kappa > 0$ , these formulas agree with the earlier results of statistical physics [25, 29, 30], and also probability theory [1, 5] in the two exactly-solvable cases where  $u_0 = \delta_0$  and  $u_0 \equiv 1$ .

*Keywords:* Stochastic partial differential equations, Lévy processes, Liapounov exponents, weak intermittence, the Burkholder–Davis–Gundy inequality.

*AMS 2000 subject classification:* Primary: 60H15; Secondary: 82B44.

*Running Title:* Intermittency and parabolic SPDEs.

---

\*Research supported in part by NSF grant DMS-0704024.

# 1 Introduction

Let  $\{\dot{w}(t, x)\}_{t \geq 0, x \in \mathbf{R}}$  denote space-time white noise, and  $\sigma : \mathbf{R} \rightarrow \mathbf{R}$  be a fixed Lipschitz function. Presently we study parabolic stochastic partial differential equations [SPDEs] of the following type:

$$\begin{cases} \partial_t u(t, x) = (\mathcal{L}u)(t, x) + \sigma(u(t, x))\dot{w}(t, x), \\ u(0, x) = u_0(x), \end{cases} \quad (1.1)$$

where  $t \geq 0$ ,  $x \in \mathbf{R}$ ,  $u_0$  is a measurable and nonnegative initial function, and  $\mathcal{L}$  is the  $L^2(\mathbf{R})$ -generator of a Lévy process  $X := \{X_t\}_{t \geq 0}$ ; and  $\mathcal{L}$  acts only on the variable  $x$ . We normalize  $X$  so that  $\mathbb{E} \exp(i\xi X_t) = \exp(-t\Psi(\xi))$  for all  $t \geq 0$  and  $\xi \in \mathbf{R}$ ;  $\mathcal{L}$  is described via its Fourier multiplier as  $\hat{\mathcal{L}}(\xi) = -\Psi(\xi)$  for all  $\xi \in \mathbf{R}$ . See the books by Bertoin [2] and Jacob [23] for pedagogic accounts.

Our principal aim is to study the mild solutions of (1.1), when they exist. At this point in time, we understand (1.1) only when its linearization with vanishing initial data has a strong solution. Together with E. Nualart [17], we have investigated precisely those linearized equations. That is,

$$\begin{cases} \partial_t u(t, x) = (\mathcal{L}u)(t, x) + \dot{w}(t, x), \\ u(0, x) = 0. \end{cases} \quad (1.2)$$

And we proved among other things that (1.2) has a strong solution if and only if Paul Lévy's symmetrization  $\bar{X}$  of the process  $X$  has local times, where

$$\bar{X}_t := X_t - X'_t \quad \text{for all } t \geq 0, \quad (1.3)$$

and  $X' := \{X'_t\}_{t \geq 0}$  is an independent copy of  $X$ . In fact, much of the local-time theory of symmetric 1-dimensional Lévy processes can be embedded within the analysis of SPDEs defined by (1.2); see [17] for details. We also proved in [17] that, as far as matters of existence and regularity are concerned, one does not encounter new phenomena if one adds to (1.2) Lipschitz-continuous additive nonlinearities. This is why we consider only multiplicative nonlinearities in (1.1).

Let  $\text{Lip}_\sigma$  denote the Lipschitz constant of  $\sigma$ , and recall that  $u_0$  is the initial data in (1.1). Here and throughout we assume, without further mention, that:

- (i)  $0 < \text{Lip}_\sigma < \infty$ , so that  $\sigma$  is [globally] Lipschitz and nontrivial; and
- (ii)  $u_0$  is bounded, nonnegative, and measurable.

Under these conditions, we prove that the SPDE (1.1) has a mild solution  $u := \{u(t, x)\}_{t \geq 0, x \in \mathbf{R}}$  that is unique up to a modification. More significantly, we show that the growth of  $t \mapsto u(t, x)$  is tied closely with the existence of  $u$ . With this aim in mind we define the *upper pth-moment Liapounov exponent*  $\bar{\gamma}(p)$  of  $u$  as

$$\bar{\gamma}(p) := \limsup_{t \rightarrow \infty} \frac{1}{t} \ln \mathbb{E}(|u(t, x)|^p) \quad \text{for all } p \in (0, \infty). \quad (1.4)$$

We say that  $u$  is *weakly intermittent*<sup>1</sup> if

$$\bar{\gamma}(2) > 0 \quad \text{and} \quad \bar{\gamma}(p) < \infty \quad \text{for all } p > 2. \quad (1.5)$$

We are interested primarily in establishing weak intermittence. However, let us mention also that weak intermittence can sometimes imply the much better-known notion of *full intermittency* [5, Definition III.1.1, p. 55]; the latter is the property that

$$p \mapsto \frac{\bar{\gamma}(p)}{p} \quad \text{is strictly increasing for all } p \geq 2. \quad (1.6)$$

Here is a brief justification: Evidently,  $\bar{\gamma}$  is convex and zero at zero, and hence  $p \mapsto \bar{\gamma}(p)/p$  is nondecreasing. Convexity implies readily that if in addition  $\bar{\gamma}(1) = 0$ , then (1.5) implies (1.6).<sup>2</sup> On the other hand, a sufficient condition for  $\bar{\gamma}(1) = 0$  is that  $u(t, x) \geq 0$  a.s. for all  $t > 0$  and  $x \in \mathbf{R}$ ; for then, (3.5) below shows immediately that  $E(|u(t, x)|) = E[u(t, x)]$  is bounded uniformly in  $t$ . We have proved the following: “*Whenever one has a comparison principle—such as that of Mueller [33] in the case that  $\mathcal{L} = \kappa \partial_{xx}$  and  $\sigma(x) = \lambda x$ —weak intermittence necessarily implies full intermittency.*”

Here, we do not pursue comparison principles. Rather, the principal goal of this note is to demonstrate that under various nearly-optimal conditions on  $\sigma$  and  $u_0$ , the solution  $u$  to (1.1) is weakly intermittent.

There is a big literature on intermittency that investigates the special case of (1.1) with  $\mathcal{L} = \kappa \partial_{xx}$  and  $\sigma(z) = \lambda z$  for constants  $\kappa > 0$  and  $\lambda \in \mathbf{R}$ ; that is the *parabolic Anderson model*. See, for example, [1, 5, 25, 29, 30, 32], together with their sizable combined references. The existing rigorous intermittency results all begin with a probabilistic formulation of (1.1) in terms of the Feynman–Kac formula. Presently, we introduce an analytic method that shows clearly that weak intermittence is connected intimately with the facts that: (i) (1.1) has a strong solution; and (ii)  $\sigma$  has linear growth, in one form or another. Our method is motivated very strongly by the theory of optimal regularity for analytic semigroups [31].

We would like to mention also that there is an impressive body of recent mathematical works on other Anderson models and  $L^p(\mathbf{P})$  intermittency, as well as almost-sure intermittency [7, 6, 8, 10, 11, 14, 16, 18, 19, 21, 20, 24, 28, 35, and their combined references].

A brief outline follows: In §2 we state the main results of the paper; these results are proved subsequently in §4, after we establish some a priori bounds in §3. Finally, we show in Appendix A that if the initial data is continuous, then the solution to (1.1) is continuous in probability, in fact continuous in  $L^p(\mathbf{P})$  for all  $p > 0$ . Consequently, if  $u_0$  is continuous, then  $u$  has a separable modification. As an immediate byproduct of our proof we find that when  $\mathcal{L}$  is the fractional Laplacian of index  $\alpha \in (1, 2]$  and  $u_0$  is continuous,  $u$  has a jointly Hölder-continuous modification (Example A.6).

<sup>1</sup>Our notion (1.5) of weak intermittence differs from that of [18, Definition 1.2].

<sup>2</sup>Inspect the proof of Theorem III.1.2 in Carmona and Molchanov [5, p. 55] for example.

## 2 Main results

We combine the existence result of [17] with a theorem of Hawkes [22] to deduce that (1.2) has a strong solution if and only if  $\Upsilon(\beta) < \infty$  for some  $\beta > 0$ , where

$$\Upsilon(\beta) := \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{d\xi}{\beta + 2\operatorname{Re} \Psi(\xi)} \quad \text{for all } \beta > 0. \quad (2.1)$$

Furthermore,  $\Upsilon(\beta)$  is finite for some  $\beta > 0$  if and only if it is finite for all  $\beta > 0$ . And under this integrability condition, (1.2) has a unique solution as well. For related results, see Brzeźniak and van Neerven [3].

Motivated by the preceding remarks, we consider only the case that the linearized equation (1.2) has a strong solution. That is, we suppose here and throughout that  $\Upsilon(\beta) < \infty$  for all  $\beta > 0$ . We might note that  $\Upsilon$  is decreasing,  $\Upsilon(\beta) > 0$  for all  $\beta > 0$ , and  $\lim_{\beta \uparrow \infty} \Upsilon(\beta) = 0$ .

Our next result establishes natural conditions for: (i) the existence and uniqueness of a solution to (1.1); and (ii)  $u$  to grow at most exponentially with a sharp exponent. It is possible to adapt the Hilbert-space methods of Peszat and Zabczyk [34] to derive existence and uniqueness. See also Da Prato [12] and Da Prato and Zabczyk [13]. Instead of following that route, we devise a method that shows very clearly that exponential growth is a consequence of the existence of a solution. Moreover, our method yields constants that will soon be shown to be essentially unimprovable.

Henceforth, by a “solution” to (1.1) we mean a mild solution  $u$  that satisfies the following:

$$\sup_{x \in \mathbf{R}} \sup_{t \in [0, T]} \mathbb{E} \left( |u(t, x)|^2 \right) < \infty \quad \text{for all } T > 0. \quad (2.2)$$

It turns out that solutions to (1.1) have better *a priori* integrability features. The following quantifies this remark.

**Theorem 2.1.** *Equation (1.1) has a solution  $u$  that is unique up to a modification. Moreover, for all even integers  $p \geq 2$ ,*

$$\bar{\gamma}(p) \leq \inf \left\{ \beta > 0 : \Upsilon \left( \frac{2\beta}{p} \right) < \frac{1}{(z_p \operatorname{Lip}_\sigma)^2} \right\} < \infty, \quad (2.3)$$

where  $z_p$  denotes the largest positive zero of the Hermite polynomial  $He_p$ .

**Remark 2.2.** We recall that  $He_p(x) = 2^{-p/2} H_p(x/2^{1/2})$  for all  $p > 0$  and  $x \in \mathbf{R}$ , where  $\exp(-2xt - t^2) = \sum_{k=0}^{\infty} H_k(x) t^k / k!$  for all  $t > 0$  and  $x \in \mathbf{R}$ . It is not hard to verify that

$$z_2 = 1 \quad \text{and} \quad z_4 = \sqrt{3 + \sqrt{6}} \approx 2.334. \quad (2.4)$$

This is valid simply because  $He_2(x) = x^2 - 1$  and  $He_4(x) = x^4 - 6x^2 + 3$ . In addition,  $z_p \sim 2p^{1/2}$  as  $p \rightarrow \infty$ , and  $\sup_{p \geq 1} (z_p / p^{1/2}) = 2$ ; see Carlen and Kree [4, Appendix].  $\square$

Before we explore the sharpness of (2.3), let us examine two cases that exhibit nonintermittence, in fact *subexponential growth*. The first concerns *subdiffusive growth*.

**Proposition 2.3.** *If  $u_0$  and  $\sigma$  are bounded and measurable, then for all integers  $p \geq 2$ ,*

$$\mathbb{E}(|u(t, x)|^p) = o\left(t^{p/2}\right) \quad \text{as } t \rightarrow \infty. \quad (2.5)$$

**Remark 2.4.** The preceding is optimal; for instance, when  $p = 2$ , the “ $o(t)$ ” cannot in general be improved to “ $o(t^\rho)$ ” for any  $\rho < 1$ . Indeed, consider the case that  $\mathcal{L} = -(-\Delta)^{\alpha/2}$  is the fractional Laplacian. It is easy to see that  $\Upsilon(\beta) < \infty$  for some  $\beta > 0$  iff  $\alpha \in (1, 2]$ . If  $0 < \inf_{z \in \mathbf{R}} |\sigma(z)| \leq \sup_{z \in \mathbf{R}} |\sigma(z)| < \infty$ , then  $\mathbb{E}(|u(t, x)|^2)$  is bounded above and below by constant multiples of  $t^{(\alpha-1)/\alpha}$ . We omit the details.  $\square$

For our second proposition we first recall the symmetrized Lévy process  $\bar{X}$  from (1.3).

**Proposition 2.5.** *If  $\bar{X}$  is transient, then for all integers  $p \geq 2$  there exists  $\delta(p) > 0$  such that  $\bar{\gamma}(p) = 0$  whenever  $\text{Lip}_\sigma < \delta(p)$ .*

**Example 2.6.** The conditions of Proposition 2.5 are not vacuous. For instance,  $\Psi(\xi) = |\xi|^\alpha + |\xi|^\rho$  is the exponent of a symmetric Lévy process  $\bar{X}$ . Moreover, if  $\alpha \in (0, 1)$  and  $\rho \in (1, 2]$ , then  $\bar{X}$  is transient and has local times.  $\square$

Our next result addresses the sharpness of (2.3), and establishes an easy-to-check sufficient criterion for  $u$  to be weakly intermittent. Throughout,  $\Upsilon^{-1}$  denotes the inverse to  $\Upsilon$  in the following sense:

$$\Upsilon^{-1}(t) := \sup \{ \beta > 0 : \Upsilon(\beta) > t \}, \quad (2.6)$$

where  $\sup \emptyset := 0$ .

**Theorem 2.7.** *If  $\inf_{z \in \mathbf{R}} u_0(z) > 0$  and  $q := \inf_{x \neq 0} |\sigma(x)/x| > 0$ , then*

$$\bar{\gamma}(2) \geq \Upsilon^{-1}\left(\frac{1}{q^2}\right) > 0. \quad (2.7)$$

Our next result is a ready corollary of Theorems 2.1 and 2.7; see Carmona and Molchanov [5, p. 59], Cranston and Molchanov [9], and Gärtner and den Hollander [18] for phenomenologically-similar results. It might help to recall (1.3).

**Corollary 2.8.** *If  $\sigma(x) := \lambda x$  and  $\inf_{x \in \mathbf{R}} u_0(x) > 0$ , then:*

1. *If  $\bar{X}$  is recurrent, then  $u$  is weakly intermittent;*
2. *If  $\bar{X}$  is transient, then  $u$  is weakly intermittent if and only if  $\Upsilon(\beta) \geq \lambda^{-2}$  for some  $\beta > 0$ ; and*
3. *In all the cases that  $u$  is weakly intermittent,  $\bar{\gamma}(2) = \Upsilon^{-1}(\lambda^{-2})$ .*

Even though Corollary 2.8 is concerned with a very special case of (1.1), that special case has a rich history. Indeed, Corollary 2.8 contains a moment analysis of the so-called *parabolic Anderson model* for  $\mathcal{L}$ . When  $\mathcal{L} = \kappa \partial_{xx}$ , that equation arises in the analysis of branching processes in random environment [5, 32]. If the spatial motion is a Lévy process with generator  $\mathcal{L}$ , then we arrive at (1.1) with  $\sigma(x) = \lambda x$ . For somewhat related—though not identical—reasons, the parabolic Anderson model also paves the way for a mathematical understanding of the so-called “KPZ equation” in dimension  $(1+1)$ . For further information see the original paper by Kardar, Parisi, and Zhang [26], Chapter 5 of Krug and Spohn [29], and the Introduction by Carmona and Molchanov [5].

**Example 2.9.** If the conditions of Corollary 2.8 hold, then the solution to (1.1) with  $\mathcal{L} = -\kappa(-\Delta)^{\alpha/2}$  is weakly intermittent with

$$\bar{\gamma}(2) = \left( \frac{\nu^\alpha \lambda^{2\alpha}}{\kappa} \right)^{1/(\alpha-1)} \quad \text{where} \quad \nu := \frac{\sec(\pi/\alpha)}{2^{1/\alpha}}. \quad (2.8)$$

Of course, we need  $\alpha \in (1, 2]$ , and this implies that  $\bar{X}$  is recurrent; see Remark 2.4. In order to derive (2.8), we first recall that  $\int_0^\infty dx/(1+x^\alpha) = (\pi/\alpha) \sec(\pi/\alpha)$ . Thus, a direct computation yields  $\Upsilon(\beta) = \nu \kappa^{-1/\alpha} \beta^{-1+(1/\alpha)}$  for all  $\beta > 0$ . Corollary 2.8, and a few more simple calculations, together imply (2.8). A similar argument shows that

$$\bar{\gamma}(p) \leq \frac{p}{2} \left( \frac{\nu^\alpha}{\kappa} (z_p \lambda)^{2\alpha} \right)^{1/(\alpha-1)} \quad \text{for all even integers } p \geq 2. \quad (2.9)$$

We can use this in conjunction with the Carlen–Kree inequality  $[z_p \leq 2\sqrt{p}]$ ; see Remark 2.2] to obtain explicit numerical bounds.  $\square$

In the special case that  $\mathcal{L} = \kappa \partial_{xx}$ , Example 2.9 tells that  $\bar{\gamma}(2) = \lambda^4/(8\kappa)$  for all  $x \in \mathbf{R}$ . This formula is anticipated by the earlier investigations of Lieb and Liniger [30] and Kardar [25, Eq. (2.9)] in statistical physics; it can also be deduced upon combining the results of Bertini and Cancrini [1], in the exact case  $u_0 \equiv 1$ , with Mueller’s comparison principle [33]. Carmona and Molchanov [5, p. 59] study a closely-related parabolic Anderson model in which  $\dot{w}(t, x)$  is white noise over  $(t, x) \in \mathbf{R}_+ \times \mathbf{Z}^d$ .

It is also easy to see that the bound furnished by (2.9) is nearly sharp in the case that  $\alpha = 2$  and  $p > 2$ . For example, (2.9) and the Carlen–Kree inequality  $[z_p \leq 2\sqrt{p}]$  together yield  $\bar{\gamma}(p) \leq \vartheta(p) := (p^3 \lambda^4)/\kappa$ , valid for all even integers  $p \geq 2$ . When  $p \geq 2$  is an arbitrary integer, the exact answer is  $\bar{\gamma}(p) = p(p^2 - 1)\lambda^4/(48\kappa)$  [1, 25, 30], and the limsup in the definition of  $\bar{\gamma}$  is a bona fide limit. Our bound  $\vartheta(p)$  agrees well with the exact answer in this special case. Indeed,

$$1 \leq \frac{\vartheta(p)}{\bar{\gamma}(p)} \leq 48 \left( 1 + \frac{1}{p^2 - 1} \right), \quad (2.10)$$

uniformly for all even integers  $p \geq 2$ , as well as all  $\lambda \in \mathbf{R}$  and  $\kappa \in (0, \infty)$ .

We close with a result that states roughly that if  $\sigma$  is asymptotically linear and  $\bar{X}$  is recurrent, then a sufficiently large initial data will ensure intermittence. More precisely, we have the following.

**Theorem 2.10.** *Suppose  $\bar{X}$  is recurrent, and  $q := \liminf_{|x| \rightarrow \infty} |\sigma(x)/x| > 0$ . Then, there exists  $\eta_0 > 0$  such that whenever  $\eta := \inf_{x \in \mathbf{R}} u_0(x) \geq \eta_0$ , the solution  $u$  is weakly intermittent.*

We believe this result presents a notable improvement on the content of Theorem 2.7 in the case that  $\bar{X}$  is recurrent.

### 3 A priori bounds

Before we prove the mathematical assertions of §2, let us develop some of the required background. Throughout we note the following elementary bound:

$$|\sigma(x)| \leq |\sigma(0)| + \text{Lip}_\sigma |x| \quad \text{for all } x \in \mathbf{R}. \quad (3.1)$$

Define  $\{\mathcal{P}_t\}_{t \geq 0}$  as the semigroup associated with  $\mathcal{L}$ . According to Lemma 8.1 of Foondun et al. [17], there exist transition densities  $\{p_t\}_{t > 0}$ , whence we have

$$(\mathcal{P}_t g)(x) = \int_{-\infty}^{\infty} p_t(y-x) g(y) dy \quad (t > 0). \quad (3.2)$$

For us, the following relation is also significant:  $(\mathcal{P}_t^* g)(x) = (\check{p}_t * g)(x)$ , where  $\mathcal{P}_t^*$  denotes the adjoint of  $\mathcal{P}_t$  in  $L^2(\mathbf{R})$ , and  $\check{p}_t(x) := p_t(-x)$ .

Consider

$$(\mathcal{G}u_0)(t, x) := (\mathcal{P}_t u_0)(x) = (\check{p}_t * u_0)(x) \quad \text{for all } t > 0 \text{ and } x \in \mathbf{R}. \quad (3.3)$$

We define also  $(\mathcal{G}u_0)(0, x) := u_0(x)$  for all  $x \in \mathbf{R}$ . The function  $v = \mathcal{G}u_0$  solves the nonrandom integro-differential equation

$$\begin{cases} \partial_t v = \mathcal{L}v & \text{on } (0, \infty) \times \mathbf{R}, \\ v(0, x) = u_0(x) & \text{for all } x \in \mathbf{R}. \end{cases} \quad (3.4)$$

Thus, we can follow the terminology and methods of Walsh [36] closely to deduce that (1.1) admits a mild solution  $u$  if and only if  $u$  is a predictable process that solves

$$u(t, x) = (\mathcal{G}u_0)(t, x) + \int_{-\infty}^{\infty} \int_0^t \sigma(u(s, y)) p_{t-s}(y-x) w(ds dy). \quad (3.5)$$

We begin by making two simple computations. The first is a basic potential-theoretic bound.

**Lemma 3.1.** *For all  $\beta > 0$ ,*

$$\sup_{t > 0} e^{-\beta t} \int_0^t \|p_s\|_{L^2(\mathbf{R})}^2 ds \leq \int_0^\infty e^{-\beta s} \|p_s\|_{L^2(\mathbf{R})}^2 ds = \Upsilon(\beta). \quad (3.6)$$

*Proof.* The inequality is obvious; we apply Plancherel's theorem to find that

$$\|p_s\|_{L^2(\mathbf{R})}^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-2s\operatorname{Re}\Psi(\xi)} d\xi \quad \text{for all } s > 0. \quad (3.7)$$

Therefore, Tonelli's theorem implies the remaining equality.  $\square$

Next we present our second elementary estimate.

**Lemma 3.2.** *For all  $a, b \in \mathbf{R}$  and  $\epsilon > 0$ ,*

$$(a + b)^2 \leq (1 + \epsilon)a^2 + (1 + \epsilon^{-1})b^2. \quad (3.8)$$

*Proof.* Define  $h(\epsilon)$  to be the upper bound of the lemma. Then,  $h : (0, \infty) \rightarrow \mathbf{R}_+$  is minimized at  $\epsilon = |b/a|$ , and the minimum value of  $h$  is  $a^2 + 2|ab| + b^2$ , which is in turn  $\geq (a + b)^2$ .  $\square$

Now we proceed to establish the remaining requires estimates.

For every positive  $t$  and all Borel sets  $A \subset \mathbf{R}$ , we set  $w_t(A) := \dot{w}([0, t] \times A)$ , and let  $\mathcal{F}_t$  denote the  $\sigma$ -algebra generated by all Wiener integrals of the form  $\int g(x) w_s(dx)$ , as the function  $g$  ranges over  $L^2(\mathbf{R})$  and the real number  $s$  ranges over  $[0, t]$ . Without loss of too much generality we may assume that the resulting filtration  $\mathcal{F} := \{\mathcal{F}_t\}_{t \geq 0}$  satisfies the usual conditions, else we enlarge each  $\mathcal{F}_t$  in the standard way. Here and throughout, a process is said to be *predictable* if it is predictable with respect to  $\mathcal{F}$ ; see also Walsh [36, p. 292].

Given a predictable random field  $f$ , we define

$$(\mathcal{A}f)(t, x) := \int_{-\infty}^{\infty} \int_0^t \sigma(f(s, y)) p_{t-s}(y - x) w(ds dy), \quad (3.9)$$

for all  $t \geq 0$  and  $x \in \mathbf{R}$ , provided that the stochastic integral exists in the sense of Walsh [36]. We also define a family of  $p$ -norms  $\{\|f\|_{p,\beta}\}_{\beta > 0}$ , one for each integer  $p \geq 2$ , via

$$\|f\|_{p,\beta} := \left\{ \sup_{t \geq 0} \sup_{x \in \mathbf{R}} e^{-\beta t} \mathbf{E}(|f(t, x)|^p) \right\}^{1/p}. \quad (3.10)$$

Variants of these norms appear in several places in the SPDE literature. See, in particular, Peszat and Zabczyk [34]. However, there is a subtle [but very important!] novelty here: The supremum is taken over all time.

Recall the definition of  $z_p$  from Theorem 2.1.

**Lemma 3.3.** *If  $f$  is predictable and  $\|f\|_{p,\beta} < \infty$  for a real  $\beta > 0$  and an even integer  $p \geq 2$ , then*

$$\|\mathcal{A}f\|_{p,\beta} \leq z_p(|\sigma(0)| + \operatorname{Lip}_\sigma \|f\|_{p,\beta}) \sqrt{\Upsilon\left(\frac{2\beta}{p}\right)}. \quad (3.11)$$



*Proof.* In his seminal 1976 paper [15], Burgess Davis found the optimal constants in the Burkholder–Davis–Gundy [BDG] inequality. In particular, Davis proved that for all  $t \geq 0$  and  $p \geq 2$ ,

$$z_p = \sup \left\{ \frac{\|N_t\|_{L^p(\mathbb{P})}}{\|\langle N, N \rangle_t\|_{L^{p/2}(\mathbb{P})}^2} : N \in \mathfrak{M}_p \right\}, \quad (3.12)$$

where  $0/0 := 0$  and  $\mathfrak{M}_p$  denotes the collection of all continuous  $L^p(\mathbb{P})$ -martingales. We apply Davis's form of the BDG inequality [*loc. cit.*], and find that

$$\begin{aligned} & \|(\mathcal{A}f)(t, x)\|_{L^p(\mathbb{P})}^p \\ & \leq z_p^p \mathbb{E} \left( \left| \int_{-\infty}^{\infty} dy \int_0^t ds |\sigma(f(s, y))|^2 |p_{t-s}(y - x)|^2 \right|^{p/2} \right). \end{aligned} \quad (3.13)$$

Since  $p/2$  is a positive integer, the preceding expectation can be written as

$$\mathbb{E} \left( \prod_{j=1}^{p/2} \int_{-\infty}^{\infty} dy_j \int_0^t ds_j |\sigma(f(s_j, y_j))|^2 |p_{t-s_j}(y_j - x)|^2 \right). \quad (3.14)$$

The generalized Hölder inequality tells us that

$$\mathbb{E} \left( \prod_{j=1}^{p/2} |\sigma(f(s_j, y_j))|^2 \right) \leq \prod_{j=1}^{p/2} \|\sigma(f(s_j, y_j))\|_{L^p(\mathbb{P})}^2. \quad (3.15)$$

Therefore, a little algebra shows us that

$$\|(\mathcal{A}f)(t, x)\|_{L^p(\mathbb{P})}^2 \leq z_p^2 \int_{-\infty}^{\infty} dy \int_0^t ds \|\sigma(f(s, y))\|_{L^p(\mathbb{P})}^2 |p_{t-s}(y - x)|^2. \quad (3.16)$$

Owing to (3.1) and Minkowski's inequality,

$$\begin{aligned} & \|(\mathcal{A}f)(t, x)\|_{L^p(\mathbb{P})}^2 \\ & \leq z_p^2 \int_{-\infty}^{\infty} dy \int_0^t ds (c_0 + c_1 \|f(s, y)\|_{L^p(\mathbb{P})})^2 |p_{t-s}(y - x)|^2, \end{aligned} \quad (3.17)$$

where  $c_0 := |\sigma(0)|$  and  $c_1 := \text{Lip}_\sigma$ , for brevity. Therefore, Lemmas 3.1 and 3.2 together imply the following bound, valid for all  $\beta > 0$ :

$$\begin{aligned} \|(\mathcal{A}f)(t, x)\|_{L^p(\mathbb{P})}^2 & \leq (1 + \epsilon^{-1}) z_p^2 c_0^2 e^{2\beta t/p} \Upsilon \left( \frac{2\beta}{p} \right) \\ & \quad + (1 + \epsilon) z_p^2 c_1^2 \int_{-\infty}^{\infty} dy \int_0^t ds \|f(s, y)\|_{L^p(\mathbb{P})}^2 |p_{t-s}(y - x)|^2. \end{aligned} \quad (3.18)$$

Because  $\|f(s, y)\|_{L^p(\mathbf{P})}^2 \leq \exp(2\beta s/p) \|f\|_{p,\beta}^2$ , it follows that

$$\begin{aligned} & \|(\mathcal{A}f)(t, x)\|_{L^p(\mathbf{P})}^2 \\ & \leq (1 + \epsilon^{-1}) z_p^2 c_0^2 e^{2\beta t/p} \Upsilon\left(\frac{2\beta}{p}\right) \\ & \quad + (1 + \epsilon) z_p^2 c_1^2 e^{2\beta t/p} \|f\|_{p,\beta}^2 \int_{-\infty}^{\infty} dy \int_0^t ds e^{-2\beta s/p} |p_s(y - x)|^2 \\ & \leq (1 + \epsilon^{-1}) z_p^2 c_0^2 e^{2\beta t/p} \Upsilon\left(\frac{2\beta}{p}\right) + (1 + \epsilon) z_p^2 c_1^2 e^{2\beta t/p} \|f\|_{p,\beta}^2 \Upsilon\left(\frac{2\beta}{p}\right). \end{aligned} \quad (3.19)$$

See Lemma 3.1 for the final inequality. We multiply both sides by  $\exp(-2\beta t/p)$  and optimize over  $t \geq 0$  and  $x \in \mathbf{R}$  to deduce the estimate

$$\|\mathcal{A}f\|_{p,\beta}^2 \leq z_p^2 \left\{ (1 + \epsilon^{-1}) |\sigma(0)|^2 + (1 + \epsilon) \text{Lip}_\sigma^2 \|f\|_{p,\beta}^2 \right\} \Upsilon\left(\frac{2\beta}{p}\right). \quad (3.20)$$

The preceding is valid for all  $\epsilon > 0$ . Now we choose

$$\epsilon := \begin{cases} |\sigma(0)|/(\text{Lip}_\sigma \|f\|_{p,\beta}) & \text{if } |\sigma(0)| \cdot \|f\|_{p,\beta} > 0, \\ 0 & \text{if } \sigma(0) = 0, \\ \infty & \text{if } \|f\|_{p,\beta} = 0, \end{cases} \quad (3.21)$$

to arrive at the statement of the lemma. Of course, “ $\epsilon = \infty$ ” means “send  $\epsilon \rightarrow \infty$ ” in the preceding.  $\square$

We plan to carry out a fixed-point argument in order to prove Theorem 2.1. The following result shows that the stochastic-integral operator  $f \mapsto \mathcal{A}f$  is a contraction on suitably-chosen spaces.

**Lemma 3.4.** *Choose and fix an even integer  $p \geq 2$ . For every  $\beta > 0$ , and all predictable random fields  $f$  and  $g$  that satisfy  $\|f\|_{p,\beta} + \|g\|_{p,\beta} < \infty$ ,*

$$\|\mathcal{A}f - \mathcal{A}g\|_{p,\beta} \leq z_p \text{Lip}_\sigma \sqrt{\Upsilon\left(\frac{2\beta}{p}\right)} \|f - g\|_{p,\beta}. \quad (3.22)$$

*Proof.* The proof is a variant of the preceding argument. Namely,

$$\begin{aligned} & \mathbf{E} (|(\mathcal{A}f)(t, x) - (\mathcal{A}g)(t, x)|^p) \\ & \leq z_p^p \mathbf{E} \left( \left| \int_{-\infty}^{\infty} dy \int_0^t ds |\sigma(f(s, y)) - \sigma(g(s, y))|^2 |p_{t-s}(y - x)|^2 \right|^{p/2} \right) \\ & \leq (z_p \text{Lip}_\sigma)^p \mathbf{E} \left( \left| \int_{-\infty}^{\infty} dy \int_0^t ds |f(s, y) - g(s, y)|^2 |p_{t-s}(y - x)|^2 \right|^{p/2} \right). \end{aligned} \quad (3.23)$$

We write the expectation as

$$\mathbf{E} \left( \prod_{j=1}^{p/2} \int_{-\infty}^{\infty} dy_j \int_0^t ds_j |f(s_j, y_j) - g(s_j, y_j)|^2 |p_{t-s_j}(y_j - x)|^2 \right), \quad (3.24)$$

and apply (3.15) to obtain the bound

$$\begin{aligned}
& \mathbb{E} (|(\mathcal{A}f)(t, x) - (\mathcal{A}g)(t, x)|^p) \\
& \leq (z_p \text{Lip}_\sigma)^p \left( \int_{-\infty}^{\infty} dy \int_0^t ds \|f(s, y) - g(s, y)\|_{L^p(\mathbb{P})}^2 |p_{t-s}(y - x)|^2 \right)^{p/2} \\
& \leq (z_p \text{Lip}_\sigma)^p \|f - g\|_{p, \beta}^p e^{\beta t} \left( \int_{-\infty}^{\infty} dy \int_0^t ds e^{-2\beta s/p} |p_s(y - x)|^2 \right)^{p/2}.
\end{aligned} \tag{3.25}$$

This has the desired effect; see Lemma 3.1.  $\square$

## 4 Proofs of the main results

*Proof of Theorem 2.1.* Define  $v_0(t, x) := u_0(x)$  for all  $(t, x) \in \mathbf{R}_+ \times \mathbf{R}$ . Since  $u_0$  is assumed to be bounded,  $\|v_0\|_{p, \beta} < \infty$  for all  $\beta > 0$  and all even integers  $p \geq 2$ . Now we iteratively set

$$v_{n+1}(t, x) := (\mathcal{A}v_n)(t, x) + (\mathcal{G}u_0)(t, x) \quad \text{for all } n \geq 0. \tag{4.1}$$

If we set  $\mathcal{A}v_{-1} := v_0$ , then thanks to Lemma 3.3, for all  $n \geq -1$ ,

$$\|\mathcal{A}v_{n+1}\|_{p, \beta} \leq z_p (|\sigma(0)| + \text{Lip}_\sigma \|\mathcal{A}v_n\|_{p, \beta}) \sqrt{\Upsilon \left( \frac{2\beta}{p} \right)}. \tag{4.2}$$

Since  $\lim_{\beta \rightarrow \infty} \Upsilon(\beta) = 0$ , we can always choose and fix  $\beta > 0$  such that

$$z_p^2 \text{Lip}_\sigma^2 \Upsilon \left( \frac{2\beta}{p} \right) < 1. \tag{4.3}$$

Given such a  $\beta$  we find, after a few lines of computation, that

$$\sup_{n \geq 0} \|\mathcal{A}v_n\|_{p, \beta} \leq \frac{z_p |\sigma(0)| \sqrt{\Upsilon(2\beta/p)}}{1 - z_p \text{Lip}_\sigma \sqrt{\Upsilon(2\beta/p)}}. \tag{4.4}$$

Because  $\mathcal{G}u_0$  is bounded uniformly by  $\sup_{z \in \mathbf{R}} u_0(z)$ , the preceding yields

$$\sup_{k \geq 1} \|v_k\|_{p, \beta} \leq \frac{z_p |\sigma(0)| \sqrt{\Upsilon(2\beta/p)}}{1 - z_p \text{Lip}_\sigma \sqrt{\Upsilon(2\beta/p)}} + \sup_{z \in \mathbf{R}} |u_0(z)|, \tag{4.5}$$

which is finite. Consequently, Lemma 3.4 assures us that all  $n \geq 1$ ,

$$\begin{aligned}
\|v_{n+1} - v_n\|_{p, \beta} &= \|\mathcal{A}v_n - \mathcal{A}v_{n-1}\|_{p, \beta} \\
&\leq z_p \text{Lip}_\sigma \sqrt{\Upsilon \left( \frac{2\beta}{p} \right)} \|v_n - v_{n-1}\|_{p, \beta}.
\end{aligned} \tag{4.6}$$

Because of (4.3), this proves the existence of a predictable random field  $u$  such that  $\lim_{n \rightarrow \infty} \|v_n - u\|_{p,\beta} = \lim_{n \rightarrow \infty} \|\mathcal{A}v_n - \mathcal{A}u\|_{p,\beta} = 0$ . Consequently,  $\|u\|_{p,\beta} < \infty$ ,  $\|u - \mathcal{A}u - \mathcal{G}u_0\|_{p,\beta} = 0$ , and

$$\mathbb{E}(|u(t, x) - (\mathcal{A}u)(t, x) - (\mathcal{G}u_0)(t, x)|^p) = 0 \quad \text{for all } (t, x) \in \mathbf{R}_+ \times \mathbf{R}. \quad (4.7)$$

These remarks prove all but one of the assertions of the theorem; we still need to establish that  $u$  is unique up to a modification. For that we follow the methods of Da Prato [12], Da Prato and Zabczyk [13], and especially Peszat and Zabczyk [34]: Suppose there are two solutions  $u$  and  $\bar{u}$  to (1.1). Define for all predictable random fields  $f$ , and  $T > 0$ ,

$$\|f\|_{2,\beta,T} := \left\{ \sup_{t \in [0,T]} \sup_{x \in \mathbf{R}} e^{-\beta t} \mathbb{E}(|f(t, x)|^2) \right\}^{1/2}. \quad (4.8)$$

Then, we can easily modify the proof of Lemma 3.4, using also the fact that  $z_2 = 1$  [Remark 2.2], to deduce that if  $u$  and  $\bar{u}$  are two solutions to (1.1), then the following holds for all  $T > 0$ :

$$\begin{aligned} \|u - \bar{u}\|_{2,\beta,T} &= \|\mathcal{A}u - \mathcal{A}\bar{u}\|_{2,\beta,T} \\ &\leq \text{Lip}_\sigma \sqrt{\Upsilon(\beta)} \|u - \bar{u}\|_{2,\beta,T}. \end{aligned} \quad (4.9)$$

Because  $\Upsilon(\beta)$  vanishes as  $\beta$  tends to infinity, this proves that  $\|u - \bar{u}\|_{2,\beta,T} = 0$  for all  $T > 0$  and all sufficiently large  $\beta > 0$ . This implies that  $u$  and  $\bar{u}$  are modifications of one another.  $\square$

*Proof of Proposition 2.3.* Because  $c_4 := \sup_{x \in \mathbf{R}} (|\sigma(x)| \vee |u_0(x)|) < \infty$ , the Burkholder–Davis–Gundy implies that

$$\|u(t, x)\|_{L^p(\mathbf{P})} \leq c_4 + c_4 z_p \left( \int_0^t \|p_s\|_{L^2(\mathbf{R})}^2 ds \right)^{1/2}. \quad (4.10)$$

Therefore, it suffices to prove that

$$\int_0^t \|p_s\|_{L^2(\mathbf{R})}^2 ds = o(t) \quad \text{as } t \rightarrow \infty. \quad (4.11)$$

The left-most term is equal to  $t \int_0^1 \|p_{st}\|_{L^2(\mathbf{R})}^2 ds$ . According to (3.7), the map  $s \mapsto \|p_s\|_{L^2(\mathbf{R})}^2$  is nonincreasing, and  $\lim_{s \rightarrow \infty} \|p_s\|_{L^2(\mathbf{R})} = 0$  by the dominated convergence theorem. Therefore, a second appeal to the dominated convergence theorem yields (4.11) and hence the theorem.  $\square$

*Proof of Theorem 2.7.* We aim to prove that

$$\int_0^\infty e^{-\beta t} \mathbb{E}(|u(t, x)|^2) dt = \infty \quad \text{provided that } \Upsilon(\beta) \geq q^{-2}. \quad (4.12)$$

This implies (2.7), as the following argument shows: Suppose, to the contrary, that  $E(|u(t, x)|^2) = O(\exp(\alpha t))$  as  $t \rightarrow \infty$ , where  $\Upsilon(\alpha) > q^{-2}$  and  $x \in \mathbf{R}$ . It follows from this that

$$\int_0^\infty e^{-\beta t} E(|u(t, x)|^2) dt \leq \text{const} \cdot \int_0^\infty e^{-(\beta-\alpha)t} dt, \quad (4.13)$$

and this is finite for every  $\beta \in (\alpha, \Upsilon^{-1}(q^{-2}))$ . Our finding contradicts (4.12), and thence follows (2.7). It remains to establish (4.12).

Let us introduce the following notation:

$$\begin{aligned} F_\beta(x) &:= \int_0^\infty e^{-\beta t} E(|u(t, x)|^2) dt \\ G_\beta(x) &:= \int_0^\infty e^{-\beta t} |(\check{p}_t * u_0)(x)|^2 dt \\ H_\beta(x) &:= \int_0^\infty e^{-\beta t} |p_t(x)|^2 dt. \end{aligned} \quad (4.14)$$

Because

$$\begin{aligned} E(|u(t, x)|^2) &= |(\check{p}_t * u_0)(x)|^2 + \int_{-\infty}^\infty dy \int_0^t ds E(|\sigma(u(s, y))|^2) |p_{t-s}(y-x)|^2, \end{aligned} \quad (4.15)$$

we may apply Laplace transforms to both sides, and then deduce that for all  $\beta > 0$  and  $x \in \mathbf{R}$ ,

$$F_\beta(x) = G_\beta(x) + \int_{-\infty}^\infty dy H_\beta(x-y) \int_0^\infty ds e^{-\beta s} E(|\sigma(u(s, y))|^2). \quad (4.16)$$

Because  $|\sigma(z)|^2 \geq q^2 |z|^2$  for all  $z \in \mathbf{R}$ , we are led to the following:

$$F_\beta(x) \geq G_\beta(x) + q^2 (F_\beta * H_\beta)(x). \quad (4.17)$$

This is a “renewal inequation,” and can be solved by standard methods. We will spell that argument out carefully, since we need an enhanced version shortly: If we define the linear operator  $\mathcal{H}$  by

$$(\mathcal{H}f)(x) := q^2 (H_\beta * f)(x), \quad (4.18)$$

then we can deduce that  $\mathcal{H}^n F_\beta - \mathcal{H}^{n+1} F_\beta \geq \mathcal{H}^n G_\beta$ , pointwise, for all integers  $n \geq 0$ . We sum this inequality from  $n = 0$  to  $n = N$  and find that

$$\begin{aligned} F_\beta(x) &\geq (\mathcal{H}^{N+1} F_\beta)(x) + \sum_{n=0}^N (\mathcal{H}^n G_\beta)(x) \\ &\geq \sum_{n=0}^N (\mathcal{H}^n G_\beta)(x). \end{aligned} \quad (4.19)$$

It follows, upon letting  $N$  tend to infinity, that

$$F_\beta(x) \geq \sum_{n=0}^{\infty} (\mathcal{H}^n G_\beta)(x). \quad (4.20)$$

If  $\eta := \inf_x u_0(x)$ , then  $(\check{p}_t * u_0)(x) \geq \eta$  pointwise, and hence  $G_\beta(x) \geq \eta^2/\beta$ . Consequently,

$$\begin{aligned} (\mathcal{H}G_\beta)(x) &\geq \frac{q^2\eta^2}{\beta} \cdot \int_{-\infty}^{\infty} H_\beta(x) dx \\ &= \frac{q^2\eta^2}{\beta} \cdot \Upsilon(\beta); \end{aligned} \quad (4.21)$$

consult Lemma 3.1 for the identity. We can iterate the preceding argument to deduce that  $F_\beta(x) \geq \eta^2\beta^{-1} \sum_{n=0}^{\infty} (q^2\Upsilon(\beta))^n$ , whence  $F_\beta(x) = \infty$  as long as  $\Upsilon(\beta) \geq q^{-2}$ . This verifies (4.12), and concludes our proof.  $\square$

*Proof of Proposition 2.5.* We recall the well-known fact that

$$\bar{X} \text{ is recurrent if and only if } \int_{-1}^1 \frac{d\xi}{\operatorname{Re} \Psi(\xi)} = \infty. \quad (4.22)$$

Otherwise,  $\bar{X}$  is transient; see Exercise V.6 of Bertoin [2, p. 152]. Because  $\Upsilon(\beta) < \infty$  for all  $\beta > 0$ , and since  $\operatorname{Re} \Psi(\xi) \geq 0$ , it is manifest that (4.22) is equivalent to the following:

$$\bar{X} \text{ is recurrent if and only if } \lim_{\beta \downarrow 0} \Upsilon(\beta) = \infty. \quad (4.23)$$

Consequently, when  $\bar{X}$  is transient,  $\sup_{\beta > 0} \Upsilon(\beta) = \lim_{\beta \downarrow 0} \Upsilon(\beta) < \infty$ , and the proposition follows immediately from Theorem 2.1. In fact, we can choose  $\delta(p)$  to be the reciprocal of  $z_p \{\sup_{\beta > 0} \Upsilon(\beta)\}^{1/2}$ .  $\square$

*Proof of Corollary 2.8.* Thanks to (4.22), when  $\bar{X}$  is recurrent, we can find  $\beta > 0$  such that  $\Upsilon(\beta) > 1/\lambda^2$ . Theorem 2.7 implies the exponential growth of  $u$ , and the formula for  $\bar{\gamma}(2)$  follows upon combining the quantitative bounds of Theorems 2.1 and 2.7. The case where  $\bar{X}$  is transient is proved similarly.  $\square$

We close the paper with the following.

*Proof of Theorem 2.10.* We modify the proof of Theorem 2.7, and point out only the requisite changes. First of all, let us note that for all  $q_0 \in (0, q)$  there exists  $A = A(q_0) \in [0, \infty)$  such that  $|\sigma(z)| \geq q_0|z|$  provided that  $|z| > A$ . Consequently, for all  $s \in \mathbf{R}_+$  and  $y \in \mathbf{R}$ ,

$$\begin{aligned} \mathbb{E} \left( |\sigma(u(s, y))|^2 \right) &\geq q_0^2 \mathbb{E} \left( |u(s, y)|^2 ; |u(s, y)| > A \right) \\ &\geq q_0^2 \mathbb{E} \left( |u(s, y)|^2 \right) - q_0^2 A^2. \end{aligned} \quad (4.24)$$

Eq. (4.15) implies that  $\mathbb{E}(|u(t, x)|^2)$  is bounded below by

$$\begin{aligned} |(\check{p}_t * u_0)(x)|^2 + q_0^2 \int_{-\infty}^{\infty} dy \int_0^t ds \mathbb{E}(|u(s, y)|^2) |p_{t-s}(y-x)|^2 \\ - q_0^2 A^2 \int_0^t \|p_s\|_{L^2(\mathbf{R})}^2 ds. \end{aligned} \quad (4.25)$$

We multiply both sides of the preceding display by  $\exp(-\beta t)$ , for a fixed  $\beta > 0$ , and integrate  $[dt]$  to find that

$$F_\beta(x) \geq G_\beta(x) + (\mathcal{H}F_\beta)(x) - \frac{q_0^2 A^2}{\beta} \Upsilon(\beta), \quad (4.26)$$

where the notation is borrowed from the proof of Theorem 2.7. We apply  $\mathcal{H}^n$  to both sides to deduce the following: For all integers  $n \geq 0$  and  $x \in \mathbf{R}$ ,

$$\begin{aligned} (\mathcal{H}^n F_\beta)(x) &\geq (\mathcal{H}^n G_\beta)(x) + (\mathcal{H}^{n+1} F_\beta)(x) - \frac{q_0^2 A^2}{\beta} \Upsilon(\beta) \cdot |q_0^2 \Upsilon(\beta)|^n \\ &\geq \frac{\eta^2}{\beta} \cdot |q_0^2 \Upsilon(\beta)|^n + (\mathcal{H}^{n+1} F_\beta)(x) - \frac{A^2}{\beta} \cdot |q_0^2 \Upsilon(\beta)|^{n+1}, \end{aligned} \quad (4.27)$$

thanks to the tautological bound  $u_0 \geq \eta$ . We collect terms to obtain the following key estimate for the present proof:

$$(\mathcal{H}^n F_\beta)(x) - (\mathcal{H}^{n+1} F_\beta)(x) \geq \frac{\eta^2 - A^2 q_0^2 \Upsilon(\beta)}{\beta} \times |q_0^2 \Upsilon(\beta)|^n, \quad (4.28)$$

valid for all integers  $n \geq 0$  and  $x \in \mathbf{R}$ . Because  $\bar{X}$  is recurrent, (4.23) ensures that we can choose  $\beta > 0$  sufficiently small that  $q_0 \Upsilon(\beta) > 1$ . Consequently,  $F_\beta(x) \equiv \infty$  as long as  $\eta$  is greater than  $A q_0 \Upsilon(\beta)$ . This proves the theorem; confer with the paragraph immediately following (4.12).  $\square$

## A Regularity

The goal of this section is to show that one can produce a nice modification of the solution to (1.1).

**Theorem A.1.** *If  $u_0$  is continuous, then the solution to (1.1) is continuous in  $L^p(\mathbf{P})$  for all  $p > 0$ . Consequently,  $u$  has a separable modification. If, in addition,  $u_0$  is uniformly continuous, then for all  $T, p > 0$ ,*

$$\lim_{\delta, \rho \downarrow 0} \sup_{0 \leq t \leq T} \sup_{x \in \mathbf{R}} \|u(t, x) - u(s, y)\|_{L^p(\mathbf{P})} = 0. \quad (\text{A.1})$$

This theorem is a ready consequence of the following series of Lemmas A.2, A.3, A.4, and A.5. We sketch [most of] the proofs because many of the methods of this section merely expand on those of the earlier sections.

Let  $\varpi$  denote the uniform modulus of continuity of  $u_0$ . That is,

$$\varpi(\delta) := \sup_{|a-b| < \delta} |u_0(a) - u_0(b)|. \quad (\text{A.2})$$

**Lemma A.2.** *If  $u_0$  is continuous, then so is  $(t, x) \mapsto (\mathcal{P}_t u_0)(x)$ . If  $u_0$  is uniformly continuous, then so is  $(t, x) \mapsto (\mathcal{P}_t u_0)(x)$ ; in fact for all  $\delta, \rho > 0$ ,*

$$\sup_{t \geq 0} \sup_{|x-z| \leq \delta} |(\mathcal{P}_t u_0)(x) - (\mathcal{P}_t u_0)(z)| \leq \varpi(\delta), \quad (\text{A.3})$$

and

$$\sup_{|t-s| < \rho} \sup_{x \in \mathbf{R}} |(\mathcal{P}_t u_0)(x) - (\mathcal{P}_s u_0)(x)| \leq \inf_{a > 0} \left[ \varpi(a) + A\rho \sup_{0 < \xi < 1/a} |\Psi(\xi)| \right], \quad (\text{A.4})$$

with  $A := 14 \sup_{z \in \mathbf{R}} u_0(z)$ .

*Proof.* We note that

$$(\mathcal{P}_t u_0)(x) - (\mathcal{P}_s u_0)(y) = \mathbb{E}(u_0(X_t + x) - u_0(X_s + y)). \quad (\text{A.5})$$

Because  $u_0$  is bounded, if it were continuous also, then  $(t, x) \mapsto (\mathcal{P}_t u_0)(x)$  is continuous by the dominated convergence theorem. Henceforth, we assume that  $u_0$  is uniformly continuous. Inequality (A.3) follows again from the dominated convergence theorem. As regards (A.4), we note that

$$\sup_{x \in \mathbf{R}} |(\mathcal{P}_t u_0)(x) - (\mathcal{P}_s u_0)(x)| \leq \mathbb{E} \left[ \varpi(|X_t - X_s|) \wedge 2 \sup_{z \in \mathbf{R}} u_0(z) \right]. \quad (\text{A.6})$$

Because  $|1 - \mathbb{E} \exp(i\xi(X_t - X_s))| \leq |t - s| \cdot |\Psi(\xi)|$ , Paul Lévy's characteristic-function inequality [27, Exercise 7.9, p. 112] shows that for all  $a > 0$ ,

$$\begin{aligned} \mathbb{P}\{|X_t - X_s| > a\} &\leq 7a \int_0^{1/a} \left| 1 - \mathbb{E} e^{i\xi(X_t - X_s)} \right| d\xi \\ &\leq 7|t - s| \sup_{0 < \xi < 1/a} |\Psi(\xi)|. \end{aligned} \quad (\text{A.7})$$

This completes our proof readily.  $\square$

**Lemma A.3.** *For all even integers  $p \geq 2$ ,  $x, z \in \mathbf{R}$ ,  $t \geq 0$ , and  $\beta > 0$ ,*

$$\begin{aligned} &\|(\mathcal{A}u)(t, x) - (\mathcal{A}u)(t, z)\|_{L^p(\mathbf{P})} \\ &\leq \left(\frac{p}{\pi}\right)^{1/2} \|\sigma \circ u\|_{p, \beta} e^{t\beta/p} \left\{ \int_{-\infty}^{\infty} \frac{1 - \cos(\xi|x - z|)}{\beta + 2\operatorname{Re} \Psi(\xi)} d\xi \right\}^{1/2}, \end{aligned} \quad (\text{A.8})$$

where  $(\sigma \circ u)(t, x) := \sigma(u(t, x))$ .

*Proof.* We follow the pattern of the proof of Lemma 3.4, and, after a few lines of estimates, deduce that

$$\begin{aligned} &\|(\mathcal{A}u)(t, x) - (\mathcal{A}u)(t, z)\|_{L^p(\mathbf{P})}^p \\ &\leq z_p^p \left| \int_{-\infty}^{\infty} dy \int_0^t ds \|\sigma(u(s, y))\|_{L^p(\mathbf{P})}^2 [p_{t-s}(y - x) - p_{t-s}(y - z)]^2 \right|^{p/2}. \end{aligned} \quad (\text{A.9})$$



Since  $\|\sigma(u(s, y))\|_{L^p(\mathbf{P})}^2 \leq \exp(2s\beta/p) \|\sigma \circ u\|_{p,\beta}^2$  for all  $\beta > 0$ , the preceding and the Carlen–Kree inequality (Remark 2.2) together yield

$$\begin{aligned} & \|(\mathcal{A}u)(t, x) - (\mathcal{A}u)(t, z)\|_{L^p(\mathbf{P})}^p \\ & \leq 2^p p^{p/2} \|\sigma \circ u\|_{p,\beta}^2 e^{\beta t} \left| \int_{-\infty}^{\infty} dy \int_0^t ds e^{-2s\beta/p} [p_s(y-x) - p_s(y-z)]^2 \right|^{p/2}. \end{aligned} \quad (\text{A.10})$$

In accord with Plancherel's theorem,

$$\begin{aligned} & \int_{-\infty}^{\infty} [p_s(y-x) - p_s(y-z)]^2 dy \\ & = \frac{1}{\pi} \int_{-\infty}^{\infty} (1 - \cos(\xi|x-z|)) e^{-2s\operatorname{Re} \Psi(\xi)} d\xi. \end{aligned} \quad (\text{A.11})$$

The lemma follows from this and a few more lines of computation.  $\square$

Choose  $x \in \mathbf{R}$  and  $0 \leq t \leq T$ . We can write

$$(\mathcal{A}u)(T, x) - (\mathcal{A}u)(t, x) = D_1 + D_2, \quad (\text{A.12})$$

where

$$\begin{aligned} D_1 &:= \int_{-\infty}^{\infty} \int_0^t \sigma(u(s, y)) [p_{T-s}(y-x) - p_{t-s}(y-x)] w(ds dy), \\ D_2 &:= \int_{-\infty}^{\infty} \int_t^T \sigma(u(s, y)) p_{T-s}(y-x) w(ds dy). \end{aligned} \quad (\text{A.13})$$

**Lemma A.4.** *For all even integers  $p \geq 2$  and  $\beta > 0$ ,*

$$\|D_1\|_{L^p(\mathbf{P})} \leq e^{\beta t/p} \left(\frac{p}{\pi}\right)^{1/2} \|\sigma \circ u\|_{p,\beta} \left( \int_{-\infty}^{\infty} \frac{|1 - e^{-(T-t)\Psi(\xi)}|^2}{(\beta/p) + \operatorname{Re} \Psi(\xi)} d\xi \right)^{1/2}. \quad (\text{A.14})$$

*Proof.* We adjust the beginning portion of the preceding proof, and after a few lines, arrive at the following:

$$\mathbb{E}(D_1^p) \leq 2^p p^{p/2} \|\sigma \circ u\|_{p,\beta}^p \left( \int_0^t e^{2\beta s/p} \|p_{T-s} - p_{t-s}\|_{L^2(\mathbf{R})}^2 ds \right)^{p/2}. \quad (\text{A.15})$$

Thanks to Plancherel's theorem,

$$\|p_{T-s} - p_{t-s}\|_{L^2(\mathbf{R})}^2 = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-2(t-s)\operatorname{Re} \Psi(\xi)} \left| 1 - e^{-(T-t)\Psi(\xi)} \right|^2 d\xi. \quad (\text{A.16})$$

Therefore, the Lebesgue integral in (A.15) is bounded above by

$$\frac{e^{\beta t}}{2\pi} \int_{-\infty}^{\infty} \frac{|1 - e^{-(T-t)\Psi(\xi)}|^2}{(2\beta/p) + 2\operatorname{Re} \Psi(\xi)} d\xi. \quad (\text{A.17})$$

Solve to finish.  $\square$

**Lemma A.5.** *For all even integers  $p \geq 2$  and  $\beta > 0$ ,*

$$\|D_2\|_{L^p(\mathbf{P})} \leq \sqrt{8p} e^{\beta T/p} \|\sigma \circ u\|_{p,\beta} \sqrt{\Upsilon\left(\frac{1}{T-t}\right)}. \quad (\text{A.18})$$

*Proof.* We adapt the proof of the preceding lemma, to the present setting, and deduce that

$$\|D_2\|_{L^p(\mathbf{P})} \leq 2\sqrt{p} \|\sigma \circ u\|_{p,\beta} e^{\beta T/p} \left( \int_0^{T-t} \|p_s\|_{L^2(\mathbf{R})}^2 ds \right)^{1/2}. \quad (\text{A.19})$$

But for all  $\rho > 0$ ,

$$\begin{aligned} \int_0^\rho \|p_s\|_{L^2(\mathbf{R})}^2 ds &= \frac{1}{2\pi} \int_{-\infty}^\infty \frac{1 - e^{-2\rho \operatorname{Re} \Psi(\xi)}}{2\operatorname{Re} \Psi(\xi)} d\xi \\ &\leq \frac{1}{\pi} \int_{-\infty}^\infty \frac{d\xi}{(1/\rho) + 2\operatorname{Re} \Psi(\xi)} \\ &= 2\Upsilon(1/\rho). \end{aligned} \quad (\text{A.20})$$

We have used the elementary fact that  $(1 - e^{-\rho\theta})/\theta \leq 2/(\rho^{-1} + \theta)$  for all  $\theta > 0$ . The lemma follows easily from these observations.  $\square$

One can often combine the preceding proof of Theorem A.1 with methods of Gaussian analysis and produce an *almost surely continuous* modification of  $u$ . We conclude this paper with an example of this method.

**Example A.6.** Suppose  $1 < \alpha \leq 2$  and  $\mathcal{L} = -\kappa(-\Delta)^{\alpha/2}$ . Suppose also that  $u_0$  is uniformly Hölder continuous; that is,  $\varpi(a) = O(a^\theta)$  as  $a \rightarrow 0^+$  for a fixed  $\theta > 0$ . We claim that in this case  $u$  has a modification that is continuous almost surely. We prove this claim by working out the estimates produced by Lemmas A.2, A.3, A.4, and A.5. Indeed, (A.3) and (A.4) together show that  $(t, x) \mapsto (\mathcal{P}_t u_0)(x)$  is uniformly jointly Hölder continuous with respective Hölder indices  $v := \theta/(\theta + \alpha)$  [for  $t$ ] and  $\theta$  [for  $x$ ]. A few more simple calculations show that: (i) The right-hand side of (A.8) is  $O(|x - z|^\mu)$  with  $\mu := \min(1/2, \alpha - 1)$ ; and (ii) the right-hand sides of (A.14) and (A.18) are both  $O((T - t)^\eta)$  with  $\eta := (\alpha - 1)/(2\alpha)$ . In other words, we can choose and fix  $\beta > 0$  that yields the following estimate: For all  $T, p > 0$  there exists  $a = a(p, T, \beta) \in (0, \infty)$  such that for all  $s, t \in [0, T]$  and  $x, y \in \mathbf{R}$ ,

$$\|u(t, x) - u(s, y)\|_{L^p(\mathbf{P})} \leq a (|t - s|^{v \wedge \eta} + |x - y|^{\theta \wedge \mu}). \quad (\text{A.21})$$

A suitable form of the Kolmogorov continuity theorem yields the desired Hölder-continuous modification.  $\square$

## References

- [1] Bertini, Lorenzo and Nicoletta Cancrini (1994). The stochastic heat equation: Feynman–Kac formula and intermittence, *J. Statist. Physics* **78**(5/6), 1377–1402.

- [2] Bertoin, Jean (1996). *Lévy Processes*, Cambridge University Press, Cambridge.
- [3] Brzeźniak, Zdzisław and Jan van Neerven (2003). Space-time regularity for linear stochastic evolution equations driven by spatially homogeneous noise, *J. Math. Kyoto Univ.* **43**(2), 261–303.
- [4] Carlen, Eric and Paul Kree (1991).  $L^p$  estimates for multiple stochastic integrals, *Ann. Probab.* **19**(1), 354–368.
- [5] Carmona, René A. and S. A. Molchanov (1994). Parabolic Anderson Problem and Intermittency, *Memoires of the Amer. Math. Soc.* **108**, American Mathematical Society, Rhode Island.
- [6] Carmona, René, Leonid Korolov, and Stanislav Molchanov (2001). Asymptotics for the almost sure Lyapunov exponent for the solution of the parabolic Anderson problem, *Random Oper. Stochastic Equations* **9**(1), 77–86.
- [7] Carmona, René A. and Frederi Viens (1998). Almost-sure exponential behavior of a stochastic Anderson model with continuous space parameter, *Stochastics* **62**, Issues 3 & 4, 251–273.
- [8] Cranston, M. and S. Molchanov (2007). Quenched to annealed transition in the parabolic Anderson problem, *Probab. Th. Rel. Fields*, **138**(1–2), 177–193.
- [9] Cranston, M. and S. Molchanov (2007). On phase transitions and limit theorems for homopolymers, in: *Probability and Mathematical Physics*, CRM Proc. Lecture Notes **42**, 97–112, American Mathematical Society, Providence, Rhode Island.
- [10] Cranston, M., T. S. Mountford, and T. Shiga (2005). Lyapounov exponent for the parabolic Anderson model with lévy noise, *Probab. Th. Rel. Fields* **132**, 321–355.
- [11] Cranston, M. and T. S. Mountford, and T. Shiga (2002). Lyapunov exponents for the parabolic Anderson model, *Acta Math. Univ. Comenian. (N.S.)* **71**(2), 163–188.
- [12] Da Prato, Giuseppe (2007). Kolmogorov equations for stochastic PDE’s with multiplicative noise, The Second Abel Symposium, 2005, pp. 235–263. In: *Stochastic Analysis and Applications*, (ed.’s: F. E. Benth, G. di Nunn, T. Lindstrøm, Tom, B. Øksendal, and T. Zhang), Springer-Verlag, Berlin.
- [13] Da Prato, Giuseppe and Jerzy Zabczyk (1992). *Stochastic Equations in Infinite Dimensions*, Cambridge University Press, Cambridge.
- [14] Dalang, Robert C. and Carl Mueller (2008). Intermittency properties in a hyperbolic Anderson problem (preprint).
- [15] Davis, Burgess (1976). On the  $L^p$  norms of stochastic integrals and other martingales, *Duke Math. J.* **43**(4), 697–704.
- [16] Florescu, Ionuț, and Frederi Viens (2006). Sharp estimation for the almost-sure Lyapunov exponent of the Anderson model in continuous space, *Probab. Th. Rel. Fields* **135**(4), 603–644.
- [17] Foondun, Mohammud, Davar Khoshnevisan, and Eulalia Nualart (2007). A local-time correspondence for stochastic partial differential equations (preprint).
- [18] Gärtner, J. and F. den Hollander, (2006). Intermittency in a catalytic random medium, *Ann. Probab.* **34**(6), 2219–2287.
- [19] Gärtner, Jürgen and Wolfgang König (2005). The parabolic Anderson model, in: *Interactive Stochastic Systems*, 153–179, Springer, Berlin.

- [20] Gärtner, J., W. König, and S. A. Molchanov (2000). Almost sure asymptotics for the continuous parabolic Anderson model, *Probab. Th. Rel. Fields* **118**(4) 547–573.
- [21] Grüninger, Gabriela, and Wolfgang König (2008). Potential confinement property of the parabolic Anderson model (preprint).
- [22] Hawkes, John (1986). Local times as stationary processes, in: *From Local Times to Global Geometry, Control and Physics* (Coventry, 1984/85), 111–120, Pitman Res. Notes Math. Ser., Longman Sci. Tech., Harlow.
- [23] Jacob, Niels (2005). *Pseudo Differential Operators and Markov Processes*, Vol. III, Imperial College Press, London.
- [24] Hofsted, Remco van der, Wolfgang König, and Peter Mörters (2006). The universality classes in the parabolic Anderson model, *Comm. Math. Phys.* **267**(2), 307–353.
- [25] Kardar, Mehran (1987). Replica Bethe ansatz studies of two-dimensional interfaces with quenched random impurities, *Nuclear Phys.* **B290**, 582–602.
- [26] Kardar, Mehran, Giorgio Parisi, and Yi-Cheng Zhang (1986). Dynamic scaling of growing interfaces, *Phys. Rev. Lett.* **56**(9), 889–892.
- [27] Khoshnevisan, Davar (2007). *Probability*, American Mathematical Society, Providence, Rhode Island.
- [28] König, Wolfgang, Hubert Lacoin, Peter Mörters, and Nadia Sidorova (2008). A two cities theorem for the parabolic Anderson model (preprint).
- [29] Krug, J. and H. Spohn (1991). Kinetic roughening of growing surfaces, in: *Solids Far From Equilibrium: Growth, Morphology, and Defects* (ed.: C. Godrèche), Cambridge University Press, Cambridge.
- [30] Lieb, Elliott H. and Werner Liniger (1963). Exact analysis of an interacting Bose gas. I. The general solution and the ground state. *Phys. Rev. (2)* **130**, 1605–1616.
- [31] Lunardi, Alessandra (1995). *Analytic Semigroups and Optimal Regularity in Parabolic Problems*, Birkhäuser-Verlag, Basel.
- [32] Molchanov, Stanislav A. (1991). Ideas in the theory of random media, *Acta Appl. Math.* **22**, 139–282.
- [33] Mueller, Carl (1991). On the support of solutions to the heat equation with noise, *Stochastics and Stochastics Rep.* **37**(4), 225–245.
- [34] Peszat, Szymon and Jerzy Zabczyk (2000). Nonlinear stochastic wave and heat equations, *Probab. Th. Rel. Fields* **116**(3), 421–443.
- [35] Shiga, Tokuzo (1997). Exponential decay rate of survival probability in a disastrous random environment, *Probab. Th. Rel. Fields* **108**(3), 417–439.
- [36] Walsh, John B. (1986). *An Introduction to Stochastic Partial Differential Equations*, in: *École d’été de probabilités de Saint-Flour, XIV—1984*, 265–439, Lecture Notes in Math., vol. 1180, Springer, Berlin.

**Mohammad Foondun & Davar Khoshnevisan**

Department of Mathematics, University of Utah, Salt Lake City, UT 84112-0090

*Emails:* mohammad@math.utah.edu & davar@math.utah.edu

*URLs:* <http://www.math.utah.edu/~mohammad> & <http://www.math.utah.edu/~davar>